**Table 1.** Observer coverage of catch landed in metric tons (mt) by non-sablefish-endorsed fixed-gear vessels by port group and year, as recorded on fish tickets.

	Longline						
	Not	Observed	Ok	Total			
		Percent of					
		unobserved		observed mt			
Port Group	mt	mt by port	mt	by port	mt		
2004	1111	group	1111	group	1111		
Neah Bay							
Astoria	1.5	100%	0	0%	1.5		
Newport							
Crescent City							
Eureka							
Fort Bragg San Francisco	10 5	1000/	_	0%	10.5		
Monterey	10.5 29.8	100% 94%	0 2	0% 6%	31.8		
Santa Barbara	49.2	97%	1.5	3%	50.7		
Los Angeles	145.5	93%	11.8	8%	157.2		
ALL PORTS	236.4	94%	15.3	6%	251.7		
2005		0.70		0,0			
Puget Sound							
Neah Bay	0.4	1000/		201	0.4		
Astoria	3.1	100%	0	0%	3.1		
Newport Coos Bay							
Crescent City	8.2	97%	0.3	3%	8.4		
Eureka							
Fort Bragg							
San Francisco	2.8	100%	0	0%	2.8		
Monterey	78.5	99%	0.9	1%	79.4		
Santa Barbara	62.7 159.6	97% 96%	1.7 7.1	3% 4%	64.3 166.6		
Los Angeles							
ALL PORTS	314.7	97%	10	3%	324.7		
January - April 2006 Puget Sound							
Neah Bay							
Astoria							
Newport							
Coos Bay							
Crescent City	4	100%	0	0%	4		
Eureka							
San Francisco	22 E	1000/	_	Ω0/	22.5		
Monterey Santa Barbara	23.5 18.5	100% 85%	0 3.4	0% 15%	23.5 21.9		
Los Angeles	38.7	89%	4.8	11%	43.5		
ALL PORTS	84.7	91%	8.1	9%	92.8		

**Table 2.** Number of observed limited-entry non-sablefish-endorsed fixed-gear trips by year and port group.

## 2004

	Longline				
	Number Percent of tr				
Port Group	of trips	in port group			
Monterey	3	5%			
Los Angeles	49	88%			
Santa Barbara	4	7%			
ALL PORTS	56	100%			

## 2005

	Longline				
	Number	Percent of trips			
Port Group	of trips	in port group			
Newport	1	3%			
Crescent City	1	3%			
Monterey	4	10%			
Los Angeles	25	63%			
Santa Barbara	9	23%			
ALL PORTS	40	100%			

January - April 2006

	Longline							
	Number	Percent of trips						
Port Group	of trips	in port group						
Los Angeles	26	79%						
Santa Barbara	7	21%						
ALL PORTS	33	100%						

**Table 3.** Number of observed limited-entry non-sablefish-endorsed fixed-gear sets by area and year.

	2004	2005	2006
Area	Longline Sets	Longline Sets	Longline Sets
North		2	
South	123	68	57
Coastwide	123	70	57

 $\textbf{Table 4.} \ \ \text{Discard rates for species from observed limited-entry non-sable fish-endorsed fixed-gear sets south of 40° 10' N. latitude by year.}$ 

	2004		2005		2006		
Species	Pounds Percent		Pounds			Pounds Percent	
Darkblotched Rockfish							
Discard							
Retained					1	100%	
Total Catch					1	100%	
Lingcod							
Discard							
Retained	21	100%			9	100%	
Total Catch	21	100%			9	100%	
Dover sole							
Discard	295	77%	391	100%	113	92%	
Retained	90	23%			10	8%	
Total Catch	385	100%	391	100%	123	100%	
Longspine Thornyhead							
Discard	559	46%	445	36%	250	20%	
Retained	653	54%	790	64%	101	80%	
Total Catch	1,212	100%	1,235	100%	1,251	100%	
Shortspine Thornyhead	-,	,.	.,		.,		
Discard	2,736	12%	648	4%	720	6%	
Retained	21,025	88%	13,845	96%	11,160	94%	
Total Catch	23,761	100%	14,493	100%	11,880	100%	
Mixed Thornyheads	20,701	10070	,	10070	11,000	10070	
Discard	2	0%	317	10%	106	8%	
Retained	6,326	100%	2,903	90%	1,151	92%	
Total Catch	6,328	100%	3,220	100%	1,257	100%	
Sablefish	0,020	10070	0,220	10070	1,207	10070	
Discard	2,309	19%	1,313	7%	299	9%	
Retained	9,867	81%	16,803	93%	3,084	91%	
Total Catch	12,176	100%	18,116	100%	3,383	100%	
Pacific Hake	12,170	10070	10,110	10070	3,303	10070	
Discard	83	78%	9	10%	40	26%	
Retained	24	22%	82	90%	112	74%	
Total Catch	106	100%	91	100%	152	100%	
Other Flatfish	100	10070		10070	102	10070	
Discard							
Retained	28	100%					
Total Catch	28	100%					
Other Nearshore Rockfish	20	10070					
Discard							
Retained					11	100%	
Total Catch					11	100%	
Other Shelf Rockfish						10070	
Discard							
Retained	1	100%	2	100%	1	100%	
Total Catch	1	100%	2	100%	1	100%	
Other Slope Rockfish	<u>'</u>	100 /6		100 /6	I	100 /6	
Discard	69	7%	6	1%	55	2%	
Retained	976	93%	787	99%	2,708	98%	
Total Catch	1,045	100%	797 792	100%	2,763	100%	
Unidentified Roundfish	1,040	100 /0	132	100 /0	2,100	100 /0	
Discard	404	24%	11	100%	7	100%	
			1.1	100%	,	10076	
Retained	1,300	76%	4.4	1000/	7	1000/	
Total Catch	1,704	100%	11	100%	7	100%	

**Table 5.** Ratio estimates and standard errors for the total bycatch of seven overfished species and lingcod per one-hundred pounds of retained groundfish from observed limited-entry non-sablefish-endorsed fixed-gear longline sets south of 40° 10' N. latitude by year.

	2004			2005			January - April 2006		
	Number			Number			Number		
Species	of sets	catch/100 lb	se	of sets	catch/100 lb	se	of sets	catch/100 lb	se
Bocaccio Rockfish	123	0.000		68	0.000		57	0.000	
Canary Rockfish	123	0.000		68	0.000		57	0.000	
Cowcod Rockfish	123	0.000		68	0.000		57	0.000	
Darkblotched Rockfish	123	0.000		68	0.000		57	0.005	0.005
Lingcod	123	0.048	0.043	68	0.000		57	0.062	0.060
Pacific Ocean Perch	123	0.000		68	0.000		57	0.000	
Widow Rockfish	123	0.000		68	0.000		57	0.000	
Yelloweye Rockfish	123	0.000		68	0.000		57	0.000	